

SUMMARY

A study was made pertaining to the variations in sensitivity to inhibitors that may be encountered in industrial deterioration problems. The results indicate that at least three types of variations are involved: first, the variation in sensitivity of different strains, secondly, the variation in sensitivity of an individual strain, and finally, the variation produced by the particular environment in which the inhibitor is studied.

REFERENCES

- BENNETT, E. O. 1956 Control of bacterial spoilage of emulsion oils. *Soap Chem. Specialties*, **32**, No. 10, 47-49; No. 11, 46-49.
- LASSUER, P. AND LAROCHE, G. 1938 Resistance de differents types dissocies a l'action de l'acide phenique. *Trav. lab. microbiol. fac. pharm. Nancy*, **11**, 39-40.
- LOCKEMANN, G. AND ULRICH, W. 1932 Adsorption and Disinfection. IV. Vertbestimmung von Phenolderivaten nach verschieden Verfahren. *Z. Hyg. Infektionskrankh.*, **113**, 475-481.
- MARION, C. 1937 Action de l'acide phenique sur les types Ra, Rb, et S de *B. aurantiacus* tingitanus Remlinger et Bailly 1935. *Bull. assoc. diplômés microbiol. fac. pharm. Nancy*, **14**, 25-37.
- PANETH, L. 1926 Über experimentelle Veränderungen der chemischen Resistenz von Bakterien. *Klin. Wochschr.*, **5**, 1603-1606.
- PIVNIK, H. AND FABIAN, F. W. 1953 Methods for testing the germicidal value of chemical compounds for disinfecting soluble oil emulsions. *Appl. Microbiol.*, **1**, 204-207.
- SNEDECOR, G. W. 1946 *Statistical methods*, 4th ed. The Iowa State College Press; Ames, Iowa.
- SZEREMI, K. 1951 La relation entre le pouvoir desinfectant et la destruction par la chaleur. *Ann. hyg. publ. ind. et sociale*, **29**, 21-34.
- VICHER, E. E., MEYER, E., AND GATHERCOAL, E. N. 1937 Phenol resistance of *Staphylococcus aureus*. *J. Am. Pharm. Assoc.*, **26**, 590-593.
- WHEELER, H. O. AND BENNETT, E. O. 1956 Bacterial inhibitors for cutting oil. *Appl. Microbiol.*, **4**, 122-126.
- WOLF, P. H. 1945 A medium containing an acid casein hydrolyzate for use in testing disinfectants. *J. Bacteriol.*, **49**, 463-472.

Physiology of Toxin Production by *Clostridium botulinum* Types A and B

II. Effect of Carbohydrate Source on Growth, Autolysis, and Toxin Production

P. F. BONVENTRE¹ AND L. L. KEMPE

Department of Bacteriology, University of Michigan, Ann Arbor, Michigan

Received for publication June 4, 1959

The kinetics of botulinum toxin production is as yet a comparatively ill-defined process. Bonventre and Kempe (unpublished observations) have recently found that the appearance of toxin in the culture filtrates of *Clostridium botulinum* types A and B is a function not of cellular growth but of the degree of autolysis. This substantiates the report by Stone (1954), Boroff (1955), Kindler *et al.* (1955), and others that autolysis is a mechanism of toxin liberation by many of the sporeforming anaerobic bacilli. This investigation concerns itself with the effects of the exogenous carbohydrate source on the growth, autolytic and toxin synthesizing processes of *C. botulinum* types A and B.

MATERIALS AND METHODS

The organisms utilized in this study are designated as *C. botulinum* type A strain JTD-IV;² *C. botulinum*

¹ Present address: Department of Microbiology, College of Medicine, University of Cincinnati, Cincinnati, Ohio.

² Obtained through the courtesy of J. T. Duff, Fort Detrick, Frederick, Maryland.

type B, strain B-201;³ *C. botulinum* type A, strain 62-A (NCA) and *Clostridium parabotulinum* type A, strain 457-A.

A complete medium consisting of 2.0 per cent tryptic digest of casein³ and 0.5 per cent autolyzed yeast extract was employed for the cultivation of the organisms. After sterilization, the carbohydrate was added in the form of a 20 per cent sterile solution to give a final concentration of 0.5 per cent. Growth was measured turbidimetrically with a Klett-Summerson photoelectric colorimeter (model 800-3)⁴ at 600 mμ.

Toxin assay. Cell-free culture fluids were obtained by centrifugation or Seitz filtration at 4 C. Owing to the lability of the toxin, certain precautions were taken during manipulation of the samples to be assayed. Dilutions were made in cold, sterile 0.5 per cent gelatin M/15 phosphate buffer at pH 6.8. Toxicity was established by intraperitoneal inoculation of the diluted

³ Sheffield Farms Company, Inc., Chemurgic Division, New York, New York.

⁴ Klett Manufacturing Company, New York, New York.

1959]

samples in
were ther
ml) accor

Effect
ments (B
ysis of C.
of glucose
thesis was
as 0.1
long enou
found the
of *C. botu*
C. botu
in the con
and also i
added. At
tions, tox
culture fil
the abser
synthesis
carbohydr
crease in
absence o
prolonged
toxin mig
intracellu
resulted i
indicated
centration
synthesis
the less to
in absenc
Growth
carbohydr
four strai
plete me
source. S

Toxin synt
B-2C

Age of Cultu
(hr)

JTD-IV

20

48

72

B-201

20

48

72

96

samples into white mice of 20 to 25 g, and toxin titers were then calculated as minimum lethal doses (MLD/ml) according to the method of Wadsworth (1947).

RESULTS AND DISCUSSION

Effect of glucose concentration. Preliminary experiments (Bonventre, 1957) showed that the rate of autolysis of *C. botulinum* was affected by the concentration of glucose in the growth medium. Maximum toxin synthesis was obtained at concentrations of glucose as low as 0.1 per cent if the cultures were incubated long enough to allow complete autolysis. It was also found that a complete medium supported the growth of *C. botulinum* in the absence of the hexose.

C. botulinum strains JTD-IV and B-201 were grown in the complete medium containing 0.5 per cent glucose and also in the same medium to which no glucose was added. After adaptation to growth under these conditions, toxin production was determined by assaying the culture filtrates. From the data in table 1 it is seen that the absence of glucose had a profound effect on the synthesis of toxin by both strains. The absence of the carbohydrate resulted in an approximate 1000-fold decrease in the toxicity of the filtrates. Autolysis in the absence of glucose did not go to completion even on prolonged incubation at 35 C. This suggested that the toxin might not have been detected if it were present intracellularly. Sonic disintegration of the cells, however, resulted in a negligible increase in the toxicity. This indicated that glucose was required, at least in low concentrations, for the synthesis of toxin as well as for the synthesis of the autolytic enzyme system. With B-201, the less toxigenic strain of the organism, toxin synthesis in absence of glucose was negligible.

Growth and toxin synthesis in the presence of various carbohydrate sources. Growth and toxin synthesis by four strains of *C. botulinum* were followed in the complete medium containing a utilizable carbohydrate source. Strains JTD-IV, 62-A, 457-A, and B-201 were

tested for their ability to ferment the carbohydrate sources added. This was determined in phenol-red broth supplemented with the test carbohydrate at a concentration of a 0.5 per cent. Large Pyrex test tubes containing 20 ml of medium were deoxygenated by boiling for 10 min, cooled, and inoculated with 1 ml of an 18-hr culture. Transfers were made every 24 hr for 3 days to adapt the organism to the carbohydrate and to dilute out any contamination by glucose originating from the initial inoculum.

After the fourth transfer, growth was measured turbidimetrically and samples were taken periodically for toxin assay. It was found that glucose and maltose supported toxin synthesis fully. Of the other carbohydrates used, only glycerol, pyruvate and ribose partially supported synthesis of toxin. The quantity of toxin produced in the presence of galactose, lactose, xylose,




Figure 1. Effect of external carbohydrate source on growth of *Clostridium botulinum* strain JTD-IV.




Figure 2. Effect of external carbohydrate source on toxin synthesis by *Clostridium botulinum* strain JTD-IV.

TABLE 1

Toxin synthesis by *Clostridium botulinum* strains JTD-IV and B-201 in the presence and in the absence of glucose

Age of Culture (hr)	Toxin Titers (MLD/ml)			
	Culture filtrate		Culture extract	
	With glucose	No glucose	With glucose	No glucose
JTD-IV				
20	5×10^3	10^2	1×10^4	1×10^3
48	1×10^5	10^3	5×10^5	1×10^3
72	2×10^5	10^3	1×10^6	1×10^3
B-201				
20	5×10^2	Neg	—	—
48	1×10^3	Neg	—	—
72	1×10^4	Neg	—	—
96	5×10^4	Neg	1×10^5	10

and inositol was no greater than in the control medium which contained no added carbohydrate.

Growth and toxin synthesis by strain JTD-IV, which may be taken as representative for all the strains tested, are presented graphically in figure 1 and 2, respectively. Although equal growth rates were observed during the exponential growth phase with all the carbohydrates, maximum growth was achieved only with glucose and maltose. The degree to which growth was repressed with the other carbon compounds when compared to glucose, however, was not sufficient to account for the extreme differences in toxin production. It can also be noted that autolysis of the culture did not go to completion unless glucose or maltose was included in the growth medium. Sonic disintegration of the intact cells, which remained after 72 hr of incubation in the media containing carbohydrates other than glucose and maltose, did not result in the increased toxicity of the supernatant fluid showing that the toxin was not present intracellularly.

Physiology of Toxin Production by *Clostridium botulinum* Types A and B

III. Effect of pH and Temperature During Incubation on Growth, Autolysis, and Toxin Production

P. F. BONVENTRE¹ AND L. L. KEMPE

Department of Bacteriology, University of Michigan, Ann Arbor, Michigan

Received for publication June 4, 1959

The ever present hazard of botulinus intoxication has stimulated continuing research by various segments of the food industries. The fact that commercially prepared foods have been only rarely incriminated in botulinum outbreaks during the past three decades is testimony to the effectiveness of the research which has been carried out to determine the proper measures for avoiding such occurrences. The literature is voluminous regarding the upper and lower limits of pH and temperature which will support the growth of *Clostridium botulinum* in various natural food substrates and artificial media. However, these investigations are for the most part concerned with an all or none phenomenon, the organism multiplies and produces toxin or it does not. Ohye and Scott (1953, 1957) did quantitative growth determinations of *C. botulinum* types A, B, and E at various temperatures but did not attempt to cor-

relate this with toxin production. Since autolysis has been shown to be correlated with the liberation of toxin by *C. botulinum* (Boroff, 1955; Kindler *et al.*, 1955; Bonventre and Kempe, 1959), it was considered that a quantitative study concerning the effects of pH and temperature incubation on the growth, autolysis, and toxin production by *C. botulinum* would be valuable from both a practical and theoretical point of view.

MATERIALS AND METHODS

C. botulinum type A strain JTD-IV² was used in this investigation. Growth was measured turbidimetrically, and toxin assayed as described previously by Bonventre and Kempe (1959).

pH of medium. The complete medium described by Bonventre and Kempe (1959) for the cultivation of all type A strains of *C. botulinum* was freshly prepared and

² Obtained through the courtesy of J. T. Duff, Fort Detrick, Frederick, Maryland.

SUMMARY

The data obtained from this survey indicate that glucose is essential for optimal toxin synthesis and that the requirement for this hexose cannot be replaced by the other energy sources tested.

REFERENCES

- BOVENTRE, P. F. 1957 Physiological basis of toxigenicity of *Clostridium botulinum* types A and B. Ph.D. Thesis, University of Michigan, Ann Arbor, Michigan.
- BOROFF, D. A. 1955 Studies of toxins of *Clostridium botulinum*. III. Relation of autolysis to toxin production. *J. Bacteriol.*, **70**, 363-367.
- KINDLER, H., MAGER, J., AND GROSSOWICZ, N. 1955 Production of toxin by resting cells of *Clostridium parabotulinum* type A. *Science*, **122**, 926-927.
- STONE, I. L. 1954 On the mode of release of tetanus toxin from the bacterial cell. *J. Bacteriol.*, **67**, 110-116.
- WADSWORTH, A. B. 1947 *Standard methods of the Division of Laboratories and Research of the New York State Department of Health*. 3rd ed. The Williams and Wilkins Co., Baltimore, Maryland.

¹ Present address: Department of Microbiology, College of Medicine, University of Cincinnati, Cincinnati, Ohio.

adjusted
of the
cooled
found
steriliz
media
10-ml
growth
culture
numbe
drogen
incuba
ture w
with a
tion du
growth
Tem
lating
10 ml
adopte
of incu
dium w
synthes
were re
the inc
low roc
in cold
and 18

Grow
JTD-IV
Optima
pH 5.5
and 8.5




Figure
strain J